

Research brief

Summary of research paper published in Nature Communications titled: Biomass burning aerosols in most climate models are too absorbing

Hunter Brown¹, Xiaohong Liu^{1,2}, Rudra Pokhrel^{1,3}, Shane Murphy¹, Zheng Lu^{1,2}, Rawad Saleh⁴, Tero Mielonen⁵, Harri Kokkola⁵, Tommi Bergman⁶, Gunnar Myhre⁷, Ragnhild B. Skeie⁷, Duncan Watson-Parris⁸, Philip Stier⁸, Ben Johnson⁹, Nicolas Bellouin¹⁰, Michael Schulz¹¹, Ville Vakkari^{12,13}, Johan Paul Beukes¹³, Pieter Gideon van Zyl¹³, Shang Liu¹⁴ and Duli Chand¹⁵

¹Department of Atmospheric Science, University of Wyoming, Laramie, WY, USA

²Department of Atmospheric Sciences, Texas A&M University, College Station, TX, USA

³Department of Physics, North Carolina A&T State University, Greensboro, NC, USA

⁴Air Quality and Climate Research Laboratory, University of Georgia, Athens, GA, USA

⁵Finnish Meteorological Institute, FI-70211 Kuopio, Finland

⁶Climate System Research, Finnish Meteorological Institute, FI-00101 Helsinki, Finland

⁷Center for International Climate and Environmental Research – Oslo (CICERO), Oslo, Norway

⁸Atmospheric, Oceanic and Planetary Physics, Department of Physics, University of Oxford, Oxford, UK

⁹Met Office, Exeter, UK

¹⁰Department of Meteorology, University of Reading, Reading, UK

¹¹Norwegian Meteorological Institute, Oslo, Norway

¹²Finnish Meteorological Institute, FI-00101 Helsinki, Finland

¹³Atmospheric Chemistry Research Group, Chemical Resource Beneficiation, North-West University, Potchefstroom, South Africa

¹⁴School of Earth and Space Sciences, University of Science and Technology of China, Hefei, China

¹⁵Atmospheric Sciences and Global Change Division, Pacific Northwest National Laboratory, Richland, WA, USA

<https://doi.org/10.17159/caj/2021/31/1.11221>

In a recent paper published in the high-impact factor journal *Nature Communications*, data collected at the Welgegund atmospheric monitoring station was one of 12 observational datasets utilised in a study to quantify the uncertainty in the representation of biomass burning (BB) aerosol composition and optical properties in climate models. Biomass burning aerosol make up a majority of primary combustion aerosol emissions (Andreae, 2019), with the main sources of global BB mass being Africa (~52%), South America (~15%), Equatorial Asia (~10%), Boreal forests (~9%), and Australia (~7%) (Van der Werf et al., 2010). The composition, size, and mixing state of BB aerosols determine the optical properties of smoke plumes in the atmosphere, which in turn is a major factor in dictating how they perturb the energy balance in the earth system. Depending on the model, the top-of-the-atmosphere BB aerosol effect can range from cooling to warming.

By relating aerosol absorption relative to extinction and carbonaceous aerosol composition from 12 observational datasets to nine state-of-the-art Earth system models and chemical transport models, varying degrees of overestimation in BB aerosol absorptivity by these models were identified. Modifications to BB aerosol refractive index, size, and mixing state was made in the Community Atmosphere Model version 5 (CAM5), which improved the model in agreement with observational

measurements. These improvements led to a global change in BB direct radiative effect of -0.07 W.m^{-2} , while regional changes of -2 W.m^{-2} in Africa, and -0.5 W.m^{-2} in South America and Temperate regions were observed. These findings suggest that current modeled BB contributes less to warming than previously thought, largely due to treatments of aerosol mixing state.

References

Andreae, M. O. Emission of trace gases and aerosols from biomass burning – an updated assessment. *Atmos. Chem. Phys.* 19, 8523–8546 (2019).

Brown H., Liu X., Pokhrel R., Murphy S., Lu Z., Saleh R., Mielonen T., Kokkola H., Bergman T., Myhre G., Skeie R.B., Watson-Paris D., Stier P., Johnson B., Bellouin N., Schulz M., Vakkari V., Beukes J.P., van Zyl P.G., Liu S. and Chand D. Biomass burning aerosols in most climate models are too absorbing. *Nat Commun* 12:277 (2021). <https://doi.org/10.1038/s41467-020-20482-9>.

Van der Werf, G. R. et al. Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009). *Atmos. Chem. Phys.* 10, 11707–11735 (2010).